

**HHS PUBLIC ACCESS**

Author manuscript

Int J Tuberc Lung Dis. Author manuscript; available in PMC 2018 July 24.

Published in final edited form as:

Int J Tuberc Lung Dis. 2013 September ; 17(9): 1139–1150. doi:10.5588/ijtld.12.0385.**Burden of tuberculosis in indigenous peoples globally: a systematic review****D. Tollefson^{*}, E. Bloss[†], A. Fanning[‡], J. T. Redd[§], K. Barker[¶], and E. McCray[†]**^{*}Rollins School of Public Health, Emory University, Atlanta, Georgia[†]US Centers for Disease Control and Prevention, Atlanta, Georgia, USA[‡]University of Alberta, Edmonton, Alberta, Canada[§]US Indian Health Service, Santa Fe, New Mexico, USA[¶]University of Toronto, Toronto, Ontario, Canada**SUMMARY****BACKGROUND**—The burden of tuberculosis (TB) in the estimated 370 million indigenous peoples worldwide is unknown.**OBJECTIVE**—To conduct a literature review to summarize the TB burden in indigenous peoples, identify gaps in current knowledge, and provide the foundation for a research agenda prioritizing indigenous health within TB control.**METHODS**—A systematic literature review identified articles published between January 1990 and November 2011 quantifying TB disease burden in indigenous populations worldwide.**RESULTS**—Among the 91 articles from 19 countries included in the review, only 56 were from outside Australia, Canada, New Zealand and the United States. The majority of the studies showed higher TB rates among indigenous groups than non-indigenous groups. Studies from the Amazon generally reported the highest TB prevalence and incidence, but select populations from South-East Asia and Africa were found to have similarly high rates of TB. In North America, the Inuit had the highest reported TB incidence (156/100 000), whereas the Metis of Canada and American Indians/Alaska Natives experienced rates of <10/100 000. New Zealand's Maori and Pacific Islanders had higher TB incidence rates than Australian Aborigines, but all were at greater risk of developing TB than non-indigenous groups.**CONCLUSION**—Where data exist, indigenous peoples were generally found to have higher rates of TB disease than non-indigenous peoples; however, this burden varied greatly. The paucity of published information on TB burden among indigenous peoples highlights the need to implement and improve TB surveillance to better measure and understand global disparities in TB rates.**Keywords**

tuberculosis; indigenous; surveillance; global

Although the estimated 370 million indigenous people worldwide¹ comprise <5% of the global population, evidence from nations where surveillance data exist suggests that the burden of tuberculosis (TB) falls heavily on indigenous peoples^{2–6} and that they are disproportionately affected.⁷ Furthermore, of the estimated 8.8 million new cases of TB occurring each year, the burden and current state of TB in the indigenous peoples inhabiting more than 70 countries globally are not known. The Strategic Framework for Action on TB Control in Indigenous Communities emphasizes the need to increase the awareness and surveillance of TB among indigenous peoples by documenting and highlighting the incidence and burden of TB in these groups.⁸

Given the lack of information about TB among the world's indigenous peoples, this systematic literature review was conducted to better understand, document and summarize the burden of TB in indigenous peoples, identify gaps in our knowledge, develop recommendations for future work and provide the foundation to set an appropriate research agenda prioritizing indigenous health within TB control.

METHODS

Search strategy

A systematic literature review using the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) framework⁹ was conducted of articles related to the epidemiology of TB in indigenous populations published during January 1990–November 2011, focusing on articles that quantified the burden of TB disease in indigenous groups. PubMed was used to search biomedical literature from Medline, journals and online books. CabDirect was used to ensure that articles published on global issues in the applied life sciences were also searched. Subject and keyword terms used in this review are listed in Table 1.

The United Nations' definition of indigenous peoples, i.e., any group of people who self-identify as indigenous to be indigenous, was used in this paper.^{1,10} For countries with disaggregated TB data, indigenous status is largely self-reported; in these countries, the names of the indigenous groups were used for the systematic search. To ensure that the search was comprehensive, names of indigenous groups from nations where disaggregated data do not exist were gathered from Bartlett et al.'s article on indigenous people and global health research,¹⁰ and from organizations representing regional indigenous issues to the United Nations Permanent Forum on Indigenous Issues (UNPFII), specifically the Indigenous Peoples of Africa Coordinating Committee,¹¹ the Russian Association of Indigenous Peoples of the Far North¹² and the Asia Indigenous Peoples Pact.¹³ In addition, country reports submitted to the UNPFII and relevant conference abstracts on indigenous health were used to ensure as broad and inclusive a search strategy as possible.

Study selection

Articles of any language were included if they provided data on the burden of TB disease in indigenous populations that were 1) collected during or after 1990 in Australia, Canada, New Zealand or the United States, or 2) collected during or after 1985 in any other country.

We allowed the five additional years of data where research was infrequent to maximize the number of indigenous groups represented in this study. Primary and secondary research articles, including published government reports, were included. The references of included articles were reviewed to identify additional papers of interest. To ensure the most comprehensive review, studies of all sizes and scope were included. Published literature reviews summarizing findings for a particular group were not included in this review but were read to identify pertinent sources. Articles with unclear methodology regarding incidence or prevalence calculations were excluded (Figure 1).

Data extraction

The World Health Organization (WHO) regions were used to organize articles based on the geographic location of the studies. The region of the Americas was further divided into North and Latin America, the latter including Mexico and the Caribbean. Two authors independently reviewed abstracts and articles for each region and came to a consensus on the inclusion or exclusion of each article. Authors extracted the data using a standardized format developed based on study objectives, a review of similar studies, and the variables needed. Authors verified the extracted data and came to a consensus when discrepancies arose.

The following variables were collected from each included study: location, including country and province or region; name of indigenous groups(s); study design; scope of study; year(s) data were collected; number of people studied; reported burden of TB (incidence or prevalence per 100 000 population); and the TB burden in a local non-indigenous population. TB incidence and/or prevalence were calculated if the number of TB cases and a population estimate were provided. If TB rates in a local non-indigenous population were not provided, the WHO-reported national TB incidence or estimated prevalence for the year data were collected was used for comparison; if the study occurred over multiple years, the WHO reported rate was chosen for the final year of data collection. From these data, a ratio was calculated to estimate the differences in TB incidence and/or prevalence in indigenous and non-indigenous groups.

RESULTS

Subject and keyword searches conducted in PubMed and CabDirect identified 534 unique articles. Four additional articles were identified from the references of papers read. Ninety-one articles related to TB in indigenous peoples in 19 countries were included in the review (Figure 2). Over three quarters of the included articles were published during 2000–2011, but the most recent available data on TB in indigenous populations varied widely between regions and countries. Twenty-seven of 35 articles from countries with disaggregated data were published after 2000; 16 of 18 studies from Latin America and 17/21 studies from South-East Asia were also published after 2000; and studies from the African, Eastern Mediterranean and European regions were largely published before 2000.

There was also substantial variation in the design and scope of the studies included in the review, specifically between countries that had disaggregated national TB surveillance data and those that did not. Many included articles were based on studies of small populations,

reporting no levels of statistical uncertainty; as such, the quality of data within the studies varied. Table 2 summarizes the methodology and scope of the included literature.

TB burden by region

African Region—Although the estimated number of indigenous people living in Africa is 14.2 million,¹⁰ only two studies from the African region fulfilled the inclusion criteria for this review. The included studies relied on small community health surveys, accompanied by medical examinations or reviews of hospital records, to assess TB status within indigenous populations.^{14,15} The available data suggest that the prevalence of TB may be substantially higher among indigenous than non-indigenous African populations. The prevalence of TB disease was found to exceed 4000/100 000 among the nomadic Fulani in Chad,¹⁵ and 1800 among the Peul and Dogon peoples in Mali (Table 3).¹⁴

The Americas: North America—The indigenous groups of North America included in the review are comprised broadly of the following: First Nations (or North American Indians), Metis and Inuit in Canada, and American Indians (AI), Alaskan Natives (AN), Native Hawaiians (NH) and Pacific Islanders (PI) in the United States. Approximately 4.4% of Canada's population reported some indigenous ancestry as of 2009,¹⁰ and in the United States, 2.3 million and 435 000 persons were identified solely as respectively AI/AN and NH/PI in 2008.⁴⁴ There are many published studies on the TB burden among indigenous groups living in the United States and Canada. Studies from these two countries predominantly included retrospective reviews based on national governmental surveillance data (Table 3).

The most recent data indicate that TB incidence continues to be substantially higher among indigenous groups across Canada and the United States than among non-indigenous groups.^{6,16,17,44–64} In the United States, AI/AN and NH/PI had 6.8 and 22.9 times greater TB incidence than Whites in 2010.^{6,16} TB incidence rates in NH/PI are higher than in AI/AN,^{6,16,44,59} but tribal-specific data, such as among the Oglala Sioux, reveals pockets where TB incidence is much higher.⁴⁷ From 2008 to 2010, TB incidence rose for both AI/AN and NH/PI, but overall TB incidence has decreased among both groups in the last two decades.⁵³

TB incidence in indigenous populations in Canada is higher than in the United States. The northern tribes of Canada are most affected by TB: in 2009, TB incidence was 155.8/100 000 among Canada's Inuit, over 150 times the rate among non-aboriginals born in Canada.¹⁷ In comparison, national TB incidence among the Metis and the First Nations peoples were respectively 7.3 and 27.4 times higher than in non-aboriginal Canadians.¹⁷

The Americas: Latin America and the Caribbean—It is estimated that Latin America is home to 450 indigenous groups comprised of 34 million people, which make up 8% of the region's total population.¹⁰ In Brazil alone, there are an estimated 220 indigenous groups composed of 450 000 people, more than 60% of whom live in the Brazilian Amazon.⁶⁵ The majority of the studies from this region included in this review were from the Brazilian Amazon,^{19,20,22,23,66–74} with fewer studies coming from Ecuador, Paraguay, Peru and Venezuela (Table 3; Appendix*).^{18,21,24–26} No studies from the Caribbean, Mexico or

Central America were found that fit the study criteria, although indigenous communities make up large parts of their populations.¹⁰

Overall, indigenous peoples across Latin America were found to have extremely high burdens of TB. Brazilian Amazonian groups were some of the most affected, with TB incidence rates often exceeding 1000/100 000, or at least 20 times higher than the incidence in the general Brazilian population.^{67,70,73} In particular, the most recent study of the Yanomami people found the incidence of TB to be 2133/100 000, or 37 times higher than in the surrounding non-indigenous population;⁷² the prevalence of TB among the Yanomami was estimated to be up to 94 times higher than in the provincial population.⁷³ Indigenous people living in Cotopaxi Province in Ecuador's highlands had TB prevalence rates reaching 6700/100 000, over 40 times Ecuador's national prevalence.²⁴ A study of the Ache of Paraguay reported a TB incidence 75 times greater than the national incidence.¹⁸

Conversely, certain indigenous groups were found to have a TB burden lower than or equal to their non-indigenous counterparts. In Peru, for example, one study found a TB incidence in indigenous peoples similar to or lower than in the general population.²¹ Within this study, many Amazonian tribes, such as the Aguaruna, Nomastshiguenga, Shawi and Shapra, had TB incidence rates of 0.4–0.9 times the national incidence.²¹ In addition, a study among indigenous children (aged <15 years) in southwestern Brazil showed that TB incidence decreased between 2001 and 2006, attributed in part to a new TB control strategy that had been implemented.²³

Eastern Mediterranean Region—This review identified one study quantifying the burden of TB among Bedouins in Israel, which found TB incidence to be 18.1/100 000 in 1987, approximately twice as high as the 1990 national Israeli TB incidence.²⁷

European Region—The indigenous peoples of Europe are generally considered those who adhere to traditional lifestyles in the Far North. Over 40 broadly defined indigenous populations are recognized in Greenland, Scandinavia and Russia, including the Inuit, the Sami, and the Nenets, Chukchi and Evenks, respectively.^{10,63} The Roma are also considered an indigenous group prominent in this region.⁷⁵

The largest and most recent studies quantifying the TB burden in Europe were conducted in Greenland,^{29,76} where 80% of persons are indigenous Inuit (Table 3).¹⁰ From 1990 to 2001, the incidence of active TB increased in this population from 85 to 185 cases/100 000.⁷⁶ Furthermore, the most recent study found that the odds of developing active TB were 15.3 times higher in indigenous groups compared to those of Danish heritage.²⁹

Outside Greenland, there was much less information from the European region. Although Scandinavia is estimated to have approximately 70 000 Sami,⁶³ no studies were found that described the burden of TB within this population, and government data for TB in areas where the Sami predominantly reside were not disaggregated.⁶³ Very little is also known about TB within the Roma population;⁷⁵ only one study conducted among Roma was

*The Appendix is available in the online version of this article at <http://www.ingentaconnect.com/content/iatld/ijtld/2013/00000017/00000009/art00005>

identified for this review, which found the Roma in Barcelona to have a TB incidence 5.3 times greater than Spain's national TB incidence.³⁰

More than 280 000 indigenous persons are estimated to live in Russia,⁶³ but studies quantifying TB incidence or prevalence within these populations were sparse and limited in scope.^{28,77–80} The data that exist show indigenous peoples to be substantially more burdened by TB than their non-indigenous counterparts. In areas of northeastern Russia, TB prevalence was found to be 10 times higher among indigenous than non-indigenous persons.⁷⁷ However, data suggest that the TB burden has been declining, narrowing the gap between indigenous and non-indigenous groups in Russia.^{28,79} No studies with data collected within the last decade were found for this review, and thus the current TB burden among the indigenous of Russia's Far North is largely unknown.

South-East Asia Region—It is estimated that three quarters of the world's indigenous people live in Asia, with over 250 indigenous groups residing in India and 60 in Thailand alone.¹⁰ Despite the large number of indigenous people in Asia, data quantifying the burden of TB were only readily available from India.

Within India, the majority of studies were conducted in the central state of Madhya Pradesh, where tribal groups account for more than 25% of the total population (Table 3).^{32–37,81–83} Data on TB burden were primarily gathered through cross-sectional surveys conducted by academic institutions and medical research centers.

In India, the majority of tribal groups demonstrated higher TB burdens than non-tribal peoples.^{32,34–37,81–83} The Saharia were the most disproportionately affected, with a TB prevalence from 5 to >100 times greater than India's national TB prevalence.^{32,37,83} The differential burden was smaller, but still substantial, between other tribal groups and the national prevalence estimates. For example, the prevalence of TB among the Bhil, Bharia and Nicobarese tribes was 1.3–8.0 times greater than the estimated national TB prevalence, while the Baiga and Jawadhu Hills tribes were found to have TB prevalence measurements 45–50% lower than national estimates.^{33,84}

Although the burden of TB was largely found to be greater in indigenous than the non-indigenous groups in India,⁸⁵ TB prevalence varied widely between groups even within the same geographic region. Within Madhya Pradesh, TB prevalence was documented to be 0.15% among the Baiga³³ and 0.43% among the Bharia,³⁴ but between 1.4% and 46% among the Saharia tribal groups.^{35–37,83} In a study conducted in Maharashtra State, active TB was identified in only six of 46 tribal groups.³⁸ However, in tribes where active TB was observed, TB prevalence reached 730/100 000, five times greater than the provincial TB prevalence.³⁸

The data also suggest that changes in the TB burden over time have been variable within tribal groups. In Tamil Nadu's Jawadhu Hills Tribes, TB prevalence decreased by four fold from 1989 to 1993.^{82,84} However, only a minimal decline in TB has been observed among the Saharia between 1991 and the present,^{36,37,82} with reports of increased burden in some areas.⁸³

Western Pacific Region—The Western Pacific is home to many indigenous peoples: the Aborigines and Torres Strait Islanders descendants in Australia; the Maori and Pacific Islanders in New Zealand; the mountainous aborigine of Taiwan; fifty-five ethnic minority groups in China; and hills peoples and other aborigine groups in the Philippines, Indonesia, Viet Nam and surrounding nations.¹⁰ However, data on TB burden were available only for Australia, New Zealand, Taiwan and, to a lesser extent, China (Figure 2). The available data indicate that TB continues to disproportionately affect the indigenous populations of the Western Pacific Region (Table 3).

The incidence of TB was lower among the indigenous populations in Australia and New Zealand than in many regions of the world, but in comparison to the non-indigenous populations, the indigenous groups were disproportionately burdened with TB. Although TB incidence has decreased among Australian Aborigines in the past years,^{4,86–90} incidence rates remain over seven times higher among Aborigines than among Australian-born non-Aborigines.⁴ In Southern Australia and the Northern Territory, Aborigines are most disproportionately affected by TB,^{89,90} and have a 13 times higher TB incidence than their non-Aborigine counterparts.⁴

In New Zealand, Pacific Islanders were consistently found to have higher rates of TB than the Maori, but both groups have substantially higher TB incidence than their non-indigenous counterparts.^{5,91–93} The most recent studies report TB incidence to be respectively 14.3 and 34.1 times higher in Maori and Pacific Islander populations than in the non-indigenous New Zealandborn.⁵ Although the most recent data suggest that TB rates have declined among the Maori and Pacific Islanders in the past 15 years,⁵ earlier data reveal lower incidence rates among the indigenous population than currently observed.^{92,93}

As was the case for Australia and New Zealand, the burden of TB in the Taiwanese indigenous people is well documented. The most recent measure found TB incidence to be 2.7 times greater for Taiwanese residing in aborigine areas than those in non-aborigine areas.⁴³ Comparing this study to earlier studies suggests that TB incidence within the indigenous population has been falling in the last 10 years, although there remains a substantial difference in TB rates between indigenous and non-indigenous peoples.^{43,94–97}

Data on TB burden for ethnic minorities in China were scarce. Although not considered indigenous peoples by the national government, many scholars assert that China's ethnic minorities fit definitions of indigenous.¹⁰ Existing data are older, with two of the three included studies published more than 20 years ago.^{98,99} In 1985, a TB incidence of 6500/100 000 was reported among the E-Lun Chun, a northeastern ethnic group;⁹⁸ otherwise no other studies of these people were found. Surveillance data from 1990 show TB prevalence to be highly variable among ethnic minorities, ranging from 47/100 000 for the Bai people to 274/100 000 for the Uyghur, prevalence rates that were respectively 35% of the estimated national prevalence and over twice the estimated national prevalence.⁹⁹ The most recent data suggest that ethnic minorities in Sichuan Province are substantially more burdened than any of the nation's ethnic groups measured in 1990.¹⁰⁰

Indigenous peoples as refugees—Circumstances have led some indigenous peoples to leave their homelands and settle in other countries, sometimes as refugees. Among such groups, data on TB burden for Tibetan and Hmong refugees were available. The incidence of TB for Tibetan refugees living in India was 3.9 times greater than India's national TB incidence, with Tibetans living in central and south India largely more affected than those in northern India.^{42,101} Outside of India, the Tibetan refugee communities screened for TB in Minnesota, USA, and Toronto, Canada, had a TB incidence of >4000/100 000.^{102,103} In comparison, reported TB incidence rates were lower among Hmong refugees, but the Hmong people still had higher TB rates than non-indigenous peoples. Hmong living as refugees in Thailand had a nine times higher TB prevalence than the Thai national rate.^{41,104} Another study found the TB incidence among Hmong in Minnesota, USA, to be 1600/100 000.¹⁰²

DISCUSSION

This is the first systematic review of the literature to examine the TB burden in indigenous populations globally. We found that indigenous peoples in high-, middle- and low-income countries continue to bear a high and disproportionate burden of TB, but this burden varies greatly between groups and regions. The most recent available data suggest that the groups most burdened by TB are located in small regions of Latin America (e.g., the Ache in Paraguay and the Yanomami in Brazil), followed closely by tribes in India (e.g., the Saharia, in Madhya Pradesh) and Africa (e.g., the Fulani, in Chad). The groups least burdened by TB live in high-income countries. Aboriginal Australians, American Indians/Alaskan Natives and the Metis of Canada were found to have the lowest incidence of TB disease among indigenous groups, but the differential TB burden between indigenous and non-indigenous populations in these industrialized nations was greater in many studies than for those in developing nations. Within these industrialized countries, Pacific Islanders, First Nations and the Canadian Inuit bear an especially disproportionate burden of TB. Select groups in Latin America and India also had a highly disproportionate burden of TB, although some tribes from these regions had minimal or no differences in disease burden. Moreover, although the available data suggest that the TB burden has decreased among some groups, such as the Australian Aborigines,^{4,86–88} Taiwanese Aborigines,^{43,95} Russia's Far North peoples,^{28,79} and the AI/AN and NH/PI in the United States,^{6,16,44,53,59} disparities in TB case rates remain. These continuing disparities highlight the importance of prioritizing indigenous health in TB control efforts. They also highlight the need to have TB control strategies that are designed and implemented by and for indigenous communities that incorporate unique indigenous perspectives and values, as proposed by the Strategic Framework for Action on Tuberculosis Control in Indigenous Communities.⁸ Such strategies should be tailored to the distinctive cultural, social and historic realities of indigenous groups to effectively address the TB burden in different regions.

This review also demonstrates the paucity of available data on the TB burden among indigenous populations outside Australia, Canada, the United States and New Zealand. Although millions of indigenous peoples live in Africa, Asia and Latin America,¹⁰ little is currently known about how TB affects these populations. Specifically, no studies were found for the tribal groups of the Caribbean, Central America or the Middle East, and TB data

were lacking for the majority of tribes in the Far North (e.g., the Sami, the Nenets), South-East Asia/Western Pacific (e.g., indigenous of Malaysia, Cambodia, Viet Nam, Indonesia and the Philippines) and Africa. Published data from these regions included in this review often originated from small studies, as opposed to publications from Australia, Canada, the United States and New Zealand, which generally used national-level surveillance data. This suggests that there is not only a shortage of data but also a paucity of high-quality data, specifically disaggregated national-level surveillance data, that allows for the estimation of TB case rates in different racial/ethnic groups. This reveals the need to increase and improve TB surveillance among indigenous populations. Where strong TB surveillance already exists (e.g., Scandinavia, India), disaggregated data are needed for routine monitoring of TB rates and the effectiveness of TB control in indigenous groups. For example, adding a single variable (e.g., race/ethnicity) to the national TB surveillance system could inform the development of evidence-based policies and practices to effectively address TB among indigenous peoples. In countries or provinces where strong TB surveillance does not yet exist, operational research assessing the burden of TB within indigenous groups is needed. Furthermore, additional research is needed to assess the extent to which drug-resistant TB, TB-HIV (human immunodeficiency virus), other comorbidities (e.g., diabetes) and the burden of TB in children affects indigenous communities.

While the primary aim of this study was to document and summarize the burden of TB in indigenous peoples, this review highlights the need to elucidate reasons for the disproportionately high TB burden among these groups. Social determinants of health are important contributors to the burden of TB among indigenous peoples, who account for approximately 15% of the world's poor¹⁰⁵ and face disparities in terms of employment, income, and access to and quality of education and health care.^{44,106} Poverty is a key obstacle in accessing health care, including prompt TB diagnosis with sensitive tools and appropriate treatment.⁵⁴ Measures that increase indigenous people's access to quality TB diagnosis and treatment are therefore needed to improve case detection and ensure improved and targeted health service delivery to marginalized indigenous groups.¹⁰⁶ Successful models of health care in indigenous peoples exist. For example, despite the structural, economic and programmatic barriers to health faced by many AI/AN communities, a study from the United States found that AI/ANs more commonly received totally directly observed therapy (DOT) and had similar treatment completion rates compared with other racial/ethnic groups.⁴⁴ The availability of DOT through the United States' Indian Health Service (IHS), tribal, and urban Indian health care facilities and other partners, may have contributed to declines in case rates. Additional research is needed to examine structural and programmatic reasons for why the TB burden is higher in many indigenous peoples—but lower in others—and to identify successful strategies for TB prevention accordingly.

Limitations

This systematic review was most limited by the dearth of published data. In some regions of the world, the most recent TB data were from more than 20 years ago, which hinders the ability to accurately assess the current burden of TB within those indigenous populations. As TB surveillance data for indigenous people are unavailable in many areas of the world, researchers calculated estimates of TB based on small studies that sometimes included fewer

than 1000 people. Although these estimates provide invaluable information, especially given the lack of information in general, the small sample sizes in many studies increase the uncertainty of disease estimates; the level of uncertainty around these measurements was unavailable in many cases. As such, differences observed between groups should be interpreted with caution. Moreover, non-standardized methodologies and wide variation in the scope of studies included in this review can limit comparability across studies. For example, comparing state or national data with data collected from smaller studies is especially challenging. Similarly, caution must be taken when comparing TB rates of indigenous groups from different years, even if they reside in the same country (e.g., Maharashtra and Madhya Tribes in India), as external factors, such as changes in TB programs, were not considered in this review. Furthermore, when comparison data were not available, published WHO TB prevalence and incidence rates were used; however, these data also have limitations for the purposes of comparison, as they are national estimates and may therefore not necessarily reflect the non-indigenous population living closest to the indigenous peoples studied. In addition, we recognize that each country with disaggregated data defines their indigenous groups differently, making direct comparisons difficult. As with any systematic review, publication bias was also a limitation.

CONCLUSIONS

This review highlights the disparate impact of TB on the world's indigenous peoples. The results highlight a need to implement and improve TB surveillance in indigenous populations globally to ensure that the burden of TB in indigenous peoples is accurately reflected. The paucity of data that exist on the epidemiology of TB within indigenous peoples, particularly in low- and middle-income countries, demonstrates the need to invest more resources to assess the current TB burden and elucidate its underlying causes. Improved TB surveillance is needed to better measure, understand and address the underlying causes of disparities in TB between indigenous and non-indigenous peoples. Ultimately, this review reveals the need to prioritize indigenous populations worldwide in TB control.

Acknowledgments

The authors thank the following people in the Division of TB Elimination, Centers for Disease Control and Prevention (CDC), Atlanta, GA, USA, for providing their support in the translation of articles: G Pinheiro, E Kurbatova and M Chen. The findings and conclusions in this report are those of the authors and do not necessarily represent an official position of the US CDC.

Conflict of interest: none declared.

References

1. United Nations Permanent Forum on Indigenous Issues. State of the world's indigenous peoples. New York, NY, USA: UN; 2012. ST/ESA/328. <http://www.unhcr.org/refworld/docid/4b6700ed2.html> [Accessed December 2012]
2. Fanning A. Tuberculosis: 1. Introduction. CMAJ. 1999; 160:837–839. [PubMed: 10189433]
3. FitzGerald JM, Wang L, Elwood RK. Tuberculosis: 13. Control of the disease among aboriginal people in Canada. CMAJ. 2000; 162:351–355. [PubMed: 10693593]


4. Barry C, Konstantinos A. Tuberculosis notifications in Australia, 2007. *Commun Dis Intell*. 2009; 33:304–315.
5. Das D, Baker M, Calder L. Tuberculosis epidemiology in New Zealand: 1995–2004. *N Z Med J*. 2006; 119:U2249. [PubMed: 17063189]
6. Centers for Disease Control and Prevention. Decrease in reported tuberculosis cases—United States, 2009. *MMWR Morb Mortal Wkly Rep*. 2010; 59:289–294. [PubMed: 20300055]
7. Hoepfner VH, Marciniuk DD. Tuberculosis in aboriginal Canadians. *Can Respir J*. 2000; 7:141–146. [PubMed: 10859400]
8. Assembly of First Nations A Global Indigenous Peoples' Initiative to Stop TB Inuit Tapiriit Kanatami; Ottawa, ON, Canada: 2009
9. Moher D, Liberati A, Tetzlaff J, Altman DG. the PRISMA Group. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: the PRISMA statement. *BMJ*. 2009; 339(b2535)
10. Bartlett JG, Madariaga-Vignudo L, O'Neil JD, Kuhnlein HV. Identifying indigenous peoples for health research in a global context: a review of perspectives and challenges. *Int J Circumpolar Health*. 2007; 66:287–307. [PubMed: 18018843]
11. Indigenous Peoples of Africa Coordinating Committee I. [Accessed December 2012] 2010 <http://www.ipacc.org.za/eng/regional.asp>
12. Russian Association of Indigenous Peoples of the Far North Moscow, Russia: RAIPON; 2006 <http://www.raipon.org/History/Peoples/tabid/310/Default.aspx> [Accessed December 2012]
13. Asia Indigenous Peoples Pact Chiang Mai, Thailand: AIPP; 2011 <http://www.aippnet.org/home/index.php> [Accessed December 2012]
14. Carta MG, Coppo P, Carpinello B, Mounkuoro PP. Mental disorders and health care seeking in Bandiagara: a community survey in the Dogon Plateau. *Soc Psychiatry Psychiatr Epidemiol*. 1997; 32:222–229. [PubMed: 9184468]
15. Schelling E, Daoud S, Daugla DM, Diallo P, Tanner M, Zinsstag J. Morbidity and nutrition patterns of three nomadic pastoralist communities of Chad. *Acta Trop*. 2005; 95:16–25. [PubMed: 15866506]
16. Centers for Disease Control and Prevention. Trends in tuberculosis—United States, 2010. *MMWR Morb Mortal Wkly Rep*. 2011; 60:333–337. [PubMed: 21430636]
17. Ellis E, , Gallant V, , Scholten D, , Dawson K, , Phipers M. Tuberculosis in Canada 2009: pre-release Ottawa, ON, Canada: Public Health Agency of Canada; 2010
18. Hurtado AM, Hill KR, Rosenblatt W, Bender J, Scharmen T. Longitudinal study of tuberculosis outcomes among immunologically naive Ache natives of Paraguay. *Am J Phys Anthropol*. 2003; 121:134–150. [PubMed: 12740957]
19. Basta PC, Coimbra CE Jr, Escobar AL, Santos RV, Alves LC, de Fonseca LS. Survey for tuberculosis in an indigenous population of Amazonia: the Surui of Rondonia, Brazil. *Trans R Soc Trop Med Hyg*. 2006; 100:579–585. [PubMed: 16274716]
20. Machado Filho AC. Incidence of tuberculosis among indigenous people in the municipality of Sao Gabriel Cachoeira. *AM Rev Soc Bras Med Trop*. 2008; 41:243–246. [Portuguese]. [PubMed: 18719802]
21. Culqui DR, Trujillo OV, Cueva N, Aylas R, Salaverry O, Bonilla C. Tuberculosis in the indigenous population of Peru, 2008. *Rev Peru Med Exp Salud Publica*. 2010; 27:8–15. [Spanish]. [PubMed: 21072444]
22. Basta PC, Coimbra CE Jr, Welch JR, Correa Alves LC, Santos RV, Bastos Camacho LA. Tuberculosis among the Xavante Indians of the Brazilian Amazon: an epidemiological and ethnographic assessment. *Ann Hum Biol*. 2010; 37:643–657. [PubMed: 20113213]
23. Marques AM, Pompilio MA, dos Santos SC, Garnes SJ, da Cunha RV. Tuberculosis among Brazilian indigenous individuals aged less than 15 years-old in State of Mato Grosso do Sul, Brazil. *Rev Soc Bras Med Trop*. 2010; 43:700–704. [Portuguese]. [PubMed: 21181027]
24. Romero-Sandoval NC, Flores-Carrera OF, Sánchez-Pérez HJ, Sánchez-Pérez I, Mateo MM. Pulmonary tuberculosis in an indigenous community in the mountains of Ecuador. *Int J Tuberc Lung Dis*. 2007; 11:550–555. [PubMed: 17439680]

25. Romero-Amaro Z, Salazar PJ, Bracho MA, Atencio TR, Romero-Gori N, Montiel UC. Prevalence of pulmonary tuberculosis in indigenous and non-indigenous patients in the state of Zulia, 1996–2005. *Kasmera*. 2008; 36:159–168.
26. Fernandez de Larrea C, Fandino C, Lopez D, et al. Tuberculosis in subjects under 15 years of age in the population of Warao in Venezuela. *Invest Clin*. 2002; 43:35–48. [Spanish]. [PubMed: 11921746]
27. Greene VW, Dolberg OT, Alkan ML, Schlaeffer FC. Tuberculosis cases in the Negev 1978–1987: ethnicity, sex, and age. *Public Health Rev*. 1992; 20:53–60. [PubMed: 1305978]
28. Kuchеров AL, Makarov VK. Antitubercular care among the Chukotka Autonomic District population. *Probl Tuberk*. 2002; (3):3–6. [Russian].
29. Ladefoged K, Rendal T, Skifte T, Andersson M, Søbørg B, Koch A. Risk factors for tuberculosis in Greenland: case-control study. *Int J Tuberc Lung Dis*. 2011; 15:44–49. [PubMed: 21276295]
30. Casals M, Pila P, Langohr K, Millet JP, Cayla JA. Incidence of infectious diseases and survival among the Roma population: a longitudinal cohort study. *Eur J Public Health*. 2012; 22:262–266. [PubMed: 21217119]
31. Datta M, Radhamani MP, Sadacharam K, et al. Survey for tuberculosis in a tribal population in North Arcot District. *Int J Tuberc Lung Dis*. 2001; 5:240–249. [PubMed: 11326823]
32. Rao VG, Bhat J, Yadav R, Gopi PG, Selvakumar N, Wares DF. Diagnosis of pulmonary tuberculosis by symptoms among tribals in central India. *Natl Med J India*. 2010; 23:372–373.
33. Yadav R, Rao VG, Bhat J, Gopi PG, Selvakumar N, Wares DF. Prevalence of pulmonary tuberculosis amongst the Baigas—a primitive tribe of Madhya Pradesh, Central India. *Indian J Tuberc*. 2010; 57:114–116. [PubMed: 21114182]
34. Rao VG, Bhat J, Yadav R, Gopi PG, Selvakumar N, Wares DF. Prevalence of pulmonary tuberculosis among the Bharia, a primitive tribe of Madhya Pradesh, central India. *Int J Tuberc Lung Dis*. 2010; 14:368–370. [PubMed: 20132630]
35. Sharma P, Jain S, Bamezai R, Tiwari P. Increased prevalence of pulmonary tuberculosis in male adults of Sahariya tribe of India: a revised survey. *Indian J Community Med*. 2010; 35:267–271. [PubMed: 20922104]
36. Rao VG, Gopi PG, Bhat J, et al. Pulmonary tuberculosis: a public health problem amongst the Saharia, a primitive tribe of Madhya Pradesh, Central India. *Int J Infect Dis*. 2010; 14:e713–716. [PubMed: 20605504]
37. Rao VG, Gopi PG, Bhat J, Yadav R, Selvakumar N, Wares DF. Selected risk factors associated with pulmonary tuberculosis among Saharia tribe of Madhya Pradesh, central India. *Eur J Public Health*. 2012; 22:271–273. [PubMed: 21324957]
38. Narang P, Tyagi NK, Mendiratta DK, Jajoo UN, Bharambhe MS, Sushila Nayar. Prevalence of sputum-positive pulmonary tuberculosis in tribal and non-tribal populations of the Ashti and Karanja tahsils in Wardha District, Maharashtra State, India. *Int J Tuberc Lung Dis*. 1999; 3:478–482. [PubMed: 10383059]
39. Murhekar MV, Kolappan C, Gopi PG, Chakraborty AK, Sehgal SC. Tuberculosis situation among tribal population of Car Nicobar, India, 15 years after intensive tuberculosis control project and implementation of a national tuberculosis programme. *Bull World Health Organ*. 2004; 82:836–843. [PubMed: 15640919]
40. Kerketta AS, Bulliyya G, Babu BV, Mohapatra SSS, Nayak RN. Health status of the elderly population among four primitive tribes of Orissa, India: a clinico-epidemiological study. *Z Gerontol Geriatr*. 2009; 42:53–59.
41. Oeltmann JE, Varma JK, Ortega L, et al. Multidrug-resistant tuberculosis outbreak among US-bound Hmong refugees, Thailand, 2005. *Emerg Infect Dis*. 2008; 14:1715–1721. [PubMed: 18976554]
42. Bhatia S, Dranyi T, Rowley D. Tuberculosis among Tibetan refugees in India. *Soc Sci Med*. 2002; 54:423–432. [PubMed: 11824918]
43. Chang Y-M, Shen C-K, Chiu C-H, Chiang H-J, Lu L-C, Liou S-H. Burden of tuberculosis among aboriginal and non-aboriginal Taiwanese, 1996–2006. *Int J Tuberc Lung Dis*. 2011; 15:471–477. [PubMed: 21396205]

44. Bloss E, Holtz TH, Jereb J, et al. Tuberculosis in indigenous peoples in the US, 2003–2008. *Public Health Rep.* 2011; 126:677–689. [PubMed: 21886328]
45. Blackwood KS, Al-Azem A, Elliott LJ, Hershfield ES, Kabani AM. Conventional and molecular epidemiology of tuberculosis in Manitoba. *BMC Infect Dis.* 2003; 3:18. [PubMed: 12917019]
46. Brancker A, Ellis E. Recent trends in tuberculosis incidence in Canada 1980–1990. *Can Commun Dis Rep.* 1992; 18:129–132. [PubMed: 1291000]
47. Breault JL, Hoffman MG. A strategy for reducing tuberculosis among Oglala Sioux Native Americans. *Am J Prev Med.* 1997; 13:182–188. [PubMed: 9181205]
48. Clark M, Riben P, Nowgesic E. The association of housing density, isolation and tuberculosis in Canadian First Nations communities. *Int J Epidemiol.* 2002; 31:940–945. [PubMed: 12435764]
49. Cook VJ, Manfreda J, Hershfield ES. Tuberculous lymphadenitis in Manitoba: incidence, clinical characteristics and treatment. *Can Respir J.* 2004; 11:279–286. [PubMed: 15254609]
50. Dyck RF, Klomp H, Marciniuk DD, et al. The relationship between diabetes and tuberculosis in Saskatchewan: comparison of registered Indians and other Saskatchewan people. *Can J Public Health.* 2007; 98:55–59. [PubMed: 17278679]
51. Enarson DA. Tuberculosis in aboriginals in Canada. *Int J Tuberc Lung Dis.* 1998; 2(Suppl 1):S16–S22. [PubMed: 9755960]
52. Gessner BD. Incidence rates, clinical features, and case identification of pediatric tuberculosis in Alaska. *Int J Tuberc Lung Dis.* 1998; 2:378–383. [PubMed: 9613633]
53. Jung RS, Bennion JR, Sorvillo F, Bellomy A. Trends in tuberculosis mortality in the United States, 1990–2006: a population-based case-control study. *Public Health Rep.* 2010; 125:389–397. [PubMed: 20433033]
54. Kunimoto D, Sutherland K, Wooldrage K, et al. Transmission characteristics of tuberculosis in the foreign-born and the Canadian-born populations of Alberta, Canada. *Int J Tuberc Lung Dis.* 2004; 8:1213–1220. [PubMed: 15527153]
55. Nguyen D, Proulx JF, Westley J, Thibert L, Dery S, Behr MA. Tuberculosis in the Inuit community of Quebec, Canada. *Am J Respir Crit Care Med.* 2003; 168:1353–1357. [PubMed: 14500266]
56. Pepperell C, Chang AH, Wobeser W, Parsonnet J, Hoepfner VH. Local epidemic history as a predictor of tuberculosis incidence in Saskatchewan aboriginal communities. *Int J Tuberc Lung Dis.* 2011; 15:899–905. [PubMed: 21682962]
57. Phipers M. Pediatric tuberculosis in Canada. *Can Commun Dis Rep.* 2003; 29:139–142. [PubMed: 12961956]
58. Phipers M, Harris T, Power C. CNS tuberculosis: a longitudinal analysis of epidemiological and clinical features. *Int J Tuberc Lung Dis.* 2006; 10:99–103. [PubMed: 16466045]
59. Schneider E. Tuberculosis among American Indians and Alaska Natives in the United States, 1993–2002. *Am J Public Health.* 2005; 95:873–880. [PubMed: 15855468]
60. Sheardown C, Phipers M. Tuberculosis in Canada, 2001. *Can Commun Dis Rep.* 2003; 29:85–90. [PubMed: 12762132]
61. Wang L, Noertjojo K, Elwood RK, FitzGerald JM. Tuberculosis among aboriginal and nonaboriginal persons in British Columbia. *Can Respir J.* 2000; 7:151–157. [PubMed: 10859401]
62. Yip D, Bhargava R, Yao Y, Sutherland K, Manfreda J, Long R. Pediatric tuberculosis in Alberta: epidemiology and case characteristics (1990–2004). *Can J Public Health.* 2007; 98:276–280. [PubMed: 17896735]
63. Young TK. Circumpolar Health Supplements Vol. 3. Oulu, Finland: International Association of Circumpolar Health Publishers; 2008 Circumpolar health indicators: sources, data and maps.
64. Centers for Disease Control and Prevention. Tuberculosis outbreak on Standing Rock Sioux Reservation—North Dakota and South Dakota, 1987–1990. *MMWR Morb Mortal Wkly Rep.* 1991; 40:204–207. [PubMed: 1900567]
65. Coimbra CE Jr, Basta PC. The burden of tuberculosis in indigenous peoples in Amazonia, Brazil. *Trans R Soc Trop Med Hyg.* 2007; 101:635–636. [PubMed: 17467759]
66. Baruzzi RG, Barros VL, Rodrigues D, Souza AL, Pagliaro H. Health and disease among Panara (Kreen-Akarore) Indians in Central Brazil after twenty-five years of contact with our world, with

- an emphasis on tuberculosis. *Cad Saude Publica*. 2001; 17:407–412. [Portuguese]. [PubMed: 11283771]
67. Basta PC, Coimbra CE Jr, Escobar AL, Santos RV. Epidemiologic aspects of tuberculosis in the Surui Indians, Brazilian Amazon. *Rev Soc Bras Med Trop*. 2004; 37:338–342. [Portuguese]. [PubMed: 15334269]
 68. Boia MN, Carvalho-Costa FA, Sodre FC, et al. Tuberculosis and intestinal parasitism among indigenous people in the Brazilian Amazon region. *Rev Saude Publica*. 2009; 43:176–178. [Portuguese]. [PubMed: 19169591]
 69. Buchillet D, Gazin P. Tuberculosis among the Amerindian population of the upper Rio Negro region. *Cad Saude Publica*. 1998; 14:181–185. [Portuguese].
 70. Escobar AL, Coimbra CE Jr, Camacho LA, Portela MC. Tuberculosis among indigenous populations in Rondonia, Amazonia, Brazil. *Cad Saude Publica*. 2001; 17:285–298. [Portuguese]. [PubMed: 11283760]
 71. Garnelo L, Brandao LC, Levino A. Dimensions and potentialities of the geographic information system on indigenous health. *Rev Saude Publica*. 2005; 39:634–640. [Portuguese]. [PubMed: 16113915]
 72. Levino A, de Oliveira RM. Tuberculosis among the Indian population in Sao Gabriel da Cachoeira, Amazonas State, Brazil. *Cad Saude Publica*. 2007; 23:1728–1732. [Portuguese]. [PubMed: 17572824]
 73. Sousa AO, Salem JJ, Lee FK, et al. An epidemic of tuberculosis with a high rate of tuberculin anergy among a population previously unexposed to tuberculosis, the Yanomami Indians of the Brazilian Amazon. *Proc Natl Acad Sci USA*. 1997; 94:13227–13232. [PubMed: 9371828]
 74. van Crevel R, van Doorninck DJ, van Ams JE, Fat HT, Vreden SG, van der Meer JM. Tuberculosis among Trio-Indians in Surinam. *Ned Tijdschr Geneesk*. 2004; 148:425–429. [Dutch]. [PubMed: 15038203]
 75. Schaaf M. Confronting a hidden disease: TB in Roma communities New York, NY, USA: Open Society Institute; 2007
 76. Thomsen VO, Lillebaek T, Stenz F. Tuberculosis in Greenland—current situation and future challenges. *Int J Circumpolar Health*. 2004; 63(Suppl 2):225–229. [PubMed: 15736657]
 77. Chernukha AD, Chernukha AA, Nikitin Iu P, Voevoda MI. Medico-social and demographic problems among ethnic groups of the Northern Asia areas. *Probl Sotsialnoi Gig Zdravookhraneniia Istor Med*. 2003; (2):16–18. [Russian].
 78. Galkin VB, Vasilev AV. Prevalence of tuberculosis and chronic non-specific lung diseases in the towns of the Far North. *Probl Tuberk*. 1994; (3):10–13. [Russian].
 79. Tyrltina MA. Social factors and their effect on the epidemiology of tuberculosis in the Far North areas. *Probl Tuberk*. 1990; (1):12–16. [Russian].
 80. Vasil'ev AV, Golubeva TM, Zolotukhin VA, Galkin VB, Lebedev VB. Medico-social aspects of epidemiology of tuberculosis in the Extreme North. *Probl Tuberk*. 1992; (7–8):4–6. [Russian].
 81. Bhat J, Rao VG, Gopi PG, et al. Prevalence of pulmonary tuberculosis amongst the tribal population of Madhya Pradesh, central India. *Int J Epidemiol*. 2009; 38:1026–1032. [PubMed: 19509182]
 82. Chakma T, Viney Rao P, Pall S, Kaushal LS, Manjula D, Tiwary RS. Survey of pulmonary tuberculosis in a primitive tribe of Madhya Pradesh. *Indian J Tuberc*. 1996; 43:85–89.
 83. Sharma PR, Tiwari PK. Health status of Sahariya tribe of central India. *Rural Remote Health*. 2007; 7:791. [PubMed: 17638510]
 84. Balasubramanian R, Sadacharam K, Selvaraj R, et al. Feasibility of involving literate tribal youths in tuberculosis case-finding in a tribal area in Tamil Nadu. *Tubercle Lung Dis*. 1995; 76:355–359.
 85. Kaulagekar A, Radkar A. Social status makes a difference: tuberculosis scenario during National Family Health Survey-2. *Indian J Tuberc*. 2007; 54:17–23. [PubMed: 17455419]
 86. Li J, Roche P, Spencer J, et al. Tuberculosis notifications in Australia, 2003. *Commun Dis Intell*. 2004; 28:464–473.
 87. Miller M, Lin M, Spencer J. Tuberculosis notifications in Australia, 2001. *Commun Dis Intell*. 2002; 26:525–536.

88. Oliver G. Tuberculosis notifications in Australia, 1994. *Commun Dis Intell.* 1996; 20:108–115.
89. Simpson G, Clark P, Knight T. Changing patterns of tuberculosis in Far North Queensland. *Med J Aust.* 2006; 184:252.
90. Simpson G, Knight T. Tuberculosis in Far North Queensland, Australia. *Int J Tuberc Lung Dis.* 1999; 3:1096–1100. [PubMed: 10599013]
91. Calder L, Priest P. One year of tuberculosis in Auckland. *N Z Med J.* 1996; 109:164–167. [PubMed: 8657368]
92. Howie S, Voss L, Baker M, Calder L, Grimwood K, Byrnes C. Tuberculosis in New Zealand, 1992–2001: a resurgence. *Arch Dis Child.* 2005; 90:1157–1161. [PubMed: 16243871]
93. van der Oest AC, Kelly P, Hood D. The changing face of tuberculosis control in a rural district of New Zealand. *Int J Tuberc Lung Dis.* 2004; 8:969–975. [PubMed: 15305479]
94. Chan PC, Huang LM, Wu YC, et al. Tuberculosis in children and adolescents, Taiwan, 1996–2003. *Emerg Infect Dis.* 2007; 13:1361–1363. [PubMed: 18252109]
95. Yu MC, Bai KJ, Chang JH, Lee CN. Tuberculosis incidence and mortality in aboriginal areas of Taiwan, 1997–2001. *J Formos Med Assoc.* 2004; 103:817–823. [PubMed: 15549148]
96. Hsueh PR, Liu YC, So J, Liu CY, Yang PC, Luh KT. *Mycobacterium tuberculosis* in Taiwan. *J Infect.* 2006; 52:77–85. [PubMed: 16216328]
97. Chang Y-M, Tsai B-Y, Wu Y-C, Yang S-Y, Chen C-H. Risk of *Mycobacterium tuberculosis* transmission in an aboriginal village, Taiwan. *Southeast Asian J Trop Med Public Health.* 2006; 37(Suppl 3):161–164. [PubMed: 17547074]
98. Hsu HJ. Incidence of tuberculosis in the hunting tribe E-Lun-Chun in Northeast China. *Pneumologie.* 1990; 44(Suppl 1):453–454. [German]. [PubMed: 2367435]
99. Wang X, You X. An analysis of the uneven prevalence of tuberculosis in various areas and regions in China. *Zhonghua Liu Xing Bing Xue Za Zhi.* 1994; 15:195–198. [Chinese]. [PubMed: 7834700]
100. Wu J, Xiong G, Feng S, et al. Study on epidemic trend and control policy of tuberculosis in Sichuan province. *Zhonghua Jie He He Hu Xi Za Zhi.* 2002; 25:12–14. [Chinese]. [PubMed: 11953091]
101. Nelson LJ, Naik Y, Tsering K, Cegielski JP. Population-based risk factors for tuberculosis and adverse outcomes among Tibetan refugees in India, 1994–1996. *Int J Tuberc Lung Dis.* 2005; 9:1018–1026. [PubMed: 16158895]
102. Truong DH, Hedemark LL, Mickman JK, Mosher LB, Dietrich SE, Lowry PW. Tuberculosis among Tibetan immigrants from India and Nepal in Minnesota, 1992–1995. *JAMA.* 1997; 277:735–738. [PubMed: 9042846]
103. Marras TK, Wilson J, Wang EE, Amending M, Yang JW. Tuberculosis among Tibetan refugee claimants in Toronto: 1998 to 2000. *Chest.* 2003; 124:915–921. [PubMed: 12970017]
104. Centers for Disease Control and Prevention. Multidrug-resistant tuberculosis in Hmong refugees resettling from Thailand into the United States, 2004–2005. *MMWR Morb Mortal Wkly Rep.* 2005; 54:741–744. [PubMed: 16079738]
105. International Fund for Agricultural Development Statistics and key facts about indigenous peoples Rome, Italy: IFAD; 2007 <http://www.ruralpovertyportal.org/web/guest/topic/statistics/tags/indigenous%20peoples> [Accessed December 2012]
106. Ring I, Brown N. The health status of indigenous peoples and others. *BMJ.* 2003; 327:404–405. [PubMed: 12933706]

**Figure 1.**

PRISMA diagram detailing paper selection process. PRISMA = Preferred Reporting Items for Systematic Reviews and Meta-Analyses.




Figure 2.

The number of studies included in review by country. Countries with data include Canada ($n = 17$), India ($n = 17$), Brazil ($n = 13$), United States ($n = 9$), Australia ($n = 6$), Taiwan ($n = 5$), Russia ($n = 5$), New Zealand ($n = 4$), China ($n = 3$), Greenland ($n = 2$), Thailand ($n = 2$), Venezuela ($n = 2$), Chad ($n = 1$), Ecuador ($n = 1$), Israel ($n = 1$), Mali ($n = 1$), Paraguay ($n = 1$), Peru ($n = 1$) and Spain ($n = 1$). Some studies in North America covered both Canada and the United States.

Table 1

Search terminology utilized in the review process

Subject search: Tuberculosis/epidemiology, Tuberculosis/ethnology, with each continental ancestry group
Broad key word: ‘Tuberculosis’ with ‘indigenous people(s)’, ‘indigenous population(s)’, ‘tribe’ or ‘tribal’, ‘aboriginal’ or ‘aborigine’
Narrow key word: ‘Tuberculosis’ and any of the following: Maori, Inuit, American Indian, Native American, Alaska Natives, First Nations, Roma, Bushmen, Herdsmen, Hill People; <i>Asia-specific</i> : Adivasi, Lahu, Akha, Mon, Lua, Mbri, Karen, Hmong, Miao, Hui (Minority AND China); <i>Africa-specific</i> : Aka, Babenjelle, Babongo, Bacwa, Bagyeli, Baka, Bakola, Bakoya, Bambuti, Batwa, Pygmy, Aasax, Akie, Aweer, Barabaig, Dahalo, Datoga, Elmolo, Hadzabe, Hadza, Maasai, Ogiek, Sandawe, Sengwer, Waata, Yaaku, Amazigh, Imazighn, Berbers, Tuareg, Afar, Aka, Babendjelle, Boranna, Dinka, Fulani, Kanuri, Karamajong, Manjo, Nuer, Peul, Pygmy, Tuareg, Tubu, Wodaabe, Bassari, Bororo, Daza, Nemadi, Ogoni, Teda, Khoekhoe, Khoikhoi, Basarwa, Khwe, Nama, San (AND Africa), Tsumkwe; <i>Far North-specific</i> : Aleut, Alutor, Chelkancy, Chukchi, Chulymcy, Chuvancy, Dolgan, Ency, Evenk, Itelmen, Kamchadal, Kereki, Kety, Khanty, Koryak, Kumandancy, Mansi, Nanaicy, Negidarcy, Nenets, Nganasan, Nivkhy, Orochi, Oroki, Saami, Sami, Selkup, Shorcy, Soioty, Tazy, Telengity, Teleuty, Tofolar, Tubolar, Tuvini-Todjin, Udege, Ukagiry, Ulchi, Veps

Table 2

Primary study methodology and scope of included studies

	<i>n</i> (%)	Study locations
Study design		
Cohort *	59 (65)	North America (<i>n</i> = 21), Latin America (<i>n</i> = 11), Eastern Mediterranean (<i>n</i> = 1), Europe (<i>n</i> = 7), South-East Asia (<i>n</i> = 2), Western Pacific (<i>n</i> = 17)
Cross-sectional	29 (32)	Africa (<i>n</i> = 2), North America (<i>n</i> = 1), Latin America (<i>n</i> = 7), South-East Asia (<i>n</i> = 19)
Case control	3 (3)	North America (<i>n</i> = 1), Europe (<i>n</i> = 1), Western Pacific (<i>n</i> = 1)
Scope		
National	29 (32)	North America (<i>n</i> = 11), Latin America (<i>n</i> = 3), Europe (<i>n</i> = 2), South-East Asia (<i>n</i> = 2), Western Pacific (<i>n</i> = 11)
Village/town/reserve	25 (27)	Africa (<i>n</i> = 2), North America (<i>n</i> = 4), Latin America (<i>n</i> = 7), Europe (<i>n</i> = 2), South-East Asia (<i>n</i> = 10)
Province/state/region	17 (19)	North America (<i>n</i> = 6), Latin America (<i>n</i> = 2), Europe (<i>n</i> = 3), South-East Asia (<i>n</i> = 1), Western Pacific (<i>n</i> = 5)
District	9 (10)	Latin America (<i>n</i> = 1), Europe (<i>n</i> = 1), South-East Asia (<i>n</i> = 6), Western Pacific (<i>n</i> = 1)
Facility	7 (8)	North America (<i>n</i> = 1), Latin America (<i>n</i> = 3), Eastern Mediterranean (<i>n</i> = 1), South-East Asia (<i>n</i> = 2)
City	4 (4)	North America (<i>n</i> = 1), Latin America (<i>n</i> = 2), Western Pacific (<i>n</i> = 1)

* Includes TB surveillance data. TB = tuberculosis.

Table 3

Summary of the most recent data on TB burden in global indigenous populations by WHO region (see Appendix for complete list of publications)

Region, indigenous group, country	Publications with the most recent data [*]	Type of study	Scope	Study period	Prevalence or incidence/100 000 [†]		Ratio (indigenous/comparative)
					Indigenous TB burden [‡]	Comparative TB burden [§]	
Africa							
Fulani, Chad	Schelling et al., 2005 ¹⁵	Cross-sectional	Village/town/reserve	1995–2000	Prevalence: 4600	Prevalence: 429	10.7
Peul and Dogon, Mali	Carta et al., 1997 ¹⁴	Cross-sectional	Village/town/reserve	1997 [¶]	Prevalence: 1800	Prevalence: 326	5.5
North America							
Alaskan Natives and American Indians, United States	CDC, 2011 ¹⁶	Cohort study	National	2010	Incidence: 6.1	Incidence: 0.9	6.8
Native Hawaiians and Pacific Islanders, [¶] United States	CDC, 2011 ¹⁶	Cohort study	National	2010	Incidence: 20.6	Incidence: 0.9	22.9
First Nations, Canada	Ellis et al., 2010 ¹⁷	Cohort study	National	2009	Incidence: 27.4	Incidence: 1.0	27.4
Inuit, Canada	Ellis et al., 2010 ¹⁷	Cohort study	National	2009	Incidence: 155.8	Incidence: 1.0	155.8
Metis, ^{¶¶} Canada	Ellis et al., 2010 ¹⁷	Cohort study	National	2009	Incidence: 7.3	Incidence: 1.0	7.3
Latin America							
Ache natives, Paraguay	Hurtado et al., 2003 ¹⁸	Cohort study	Facility	1987–2002	Incidence: 3700	Incidence: 49	75.5
Anazonian tribes, Rondonia or Amazonas State, Brazil	Machado Filho, 2008 ²⁰ Basta et al., 2006 ¹⁹	Cross-sectional, cohort study	Facility, city	2003	Incidence: 284 Prevalence: 815.2	Incidence: 55 Prevalence: 37.5	5.2 21.7
Anazonian tribes, Peru	Culqui et al., 2010 ²¹	Cohort study	National	2008	Incidence: 42.3–400.9	Incidence: 106	0.4–3.8
Aymara (Andes), Peru	Culqui et al., 2010 ²¹	Cohort study	National	2008	Incidence: 175.1	Incidence: 106	1.7
Brazilian Native Indians, Mato Grosso do Sul, Brazil	Basta et al., 2010 ²²	Cross-sectional, cohort study	Village/town/reserve	2006, 1999–2004	Prevalence: 210.8 Incidence: 1289.6	Prevalence: 55.0 Incidence: 48.5	3.8 26.6
Chine, Ecuador	Romero-Sandoval et al., 2007 ²⁴	Cross-sectional	Village/town/reserve	2001	Prevalence: 6700	Prevalence: 161	41.6
Quechua (Andes), Peru	Culqui et al., 2010 ²¹	Cohort study	National	2008	Incidence: 103.7	Incidence: 106	1.0
Zulu State indigenous, Venezuela	Romero-Amaro et al., 2008 ²⁵	Cohort study	Province/state/region	1996–2005	52% of TB cases occurred among the indigenous	Incidence: 23.2	>1
Warao people, Venezuela	Fernandez de Larrea et al., 2002 ²⁶	Cross-sectional	Village/town/reserve	1999	Prevalence: 320.0	Prevalence: 48.0	6.7
Eastern Mediterranean							
Bedouins	Greene et al., 1992 ²⁷	Cohort study	Facility	1987	Incidence: 18.1	Incidence: 8.7	2.1
Europe							
Russian indigenous, Russia	Kucherov and Makarov, 2002 ²⁸	Cohort study	District	1999	Incidence: 85.2 Prevalence: 608	Incidence: 37.6 Prevalence: 193.3	2.3 3.1
Inuit, Greenland	Thomsen et al., 2004 ⁷⁶	Cohort study	National	2001	Prevalence: 185	Prevalence: 13.0	14.2
Roma, Spain	Casals et al., 2011 ³⁰	Cohort study	Village/town/reserve	1985–2008	Incidence: 90.9	Incidence: 17.0	5.3

Region, indigenous group, country	Publications with the most recent data [*]	Type of study	Scope	Study period	Prevalence or incidence/100 000 [†]	
					Indigenous TB burden [‡]	Comparative TB burden [§] Ratio (indigenous/comparative)
South-East Asia						
Jawadhu Hills tribes, India	Balaubramanian et al., 1995 ⁸⁴	Cross-sectional	Village/town/reserve	1992–1993	Prevalence: 209	Prevalence: 461 0.5
Madhya Pradesh tribals (including Baiga, Bhil, Bharia, Saharia), India	Rao et al., 2010 ³² Yadav et al., 2010 ³³ Rao et al., 2010 ³⁴ Bhat et al., 2009 ⁸¹ Rao et al., 2010 ³⁶ Rao et al., 2012 ³⁷	Cross-sectional	District, village/town/reserve	2007–2008	Prevalence: 146–1518	Prevalence: 294 0.5–5.2
Maharashtra tribes, India	Narang et al., 1999 ³⁸	Cross-sectional	District	1989–1990	Prevalence: 133	Prevalence: 144 0.9
Nicobarese, India	Murhekar et al., 2004 ³⁹	Cross-sectional	Province/state/region	2001–2002	Prevalence: 735.3	Prevalence: 436 1.7
Forest Peoples, Orissa, India	Kerketta et al., 2009 ⁴⁰	Cross-sectional	Village/town/reserve	2009 ^{¶¶}	Prevalence: 4808	Prevalence: 275 17.5
Hmong, Thailand	Ootmann et al., 2008 ⁴¹	Cross-sectional	Village/town/reserve	2004–2005	Prevalence: 1760 ^{¶¶}	Prevalence: 193 9.1
Tibetans, India	Bhatia et al., 2002 ⁴²	Cross-sectional	Village/town/reserve	1996	Incidence: 770	Incidence: 216 3.6
Western Pacific						
Aboriginal/Torres Strait Islanders, Australia	Barry and Konstantinos, 2009 ⁴	Cohort study	National	2007	Incidence: 6.6	Incidence: 0.9 7.3
Māori, New Zealand	Das et al., 2006 ⁵	Cohort study	National	2000–2004	Incidence: 20.46	Incidence: 1.43 14.3
Pacific Islanders, ^{†††} New Zealand	Das et al., 2006 ⁵	Cohort study	National	2000–2004	Incidence: 48.79	Incidence: 1.43 34.1
Taiwanese aboriginals, Taiwan	Chang et al., 2011 ⁴³	Cohort study	National	1996–2006	Incidence: 176	Incidence: 65 2.7
Ethnic minorities, Sichuan, China	Wu et al. 2002 ¹⁰⁰	Cohort study	Province/state/region	2000	Prevalence: 395	Prevalence: 149 (average) 2.7

^{*}When more than one study on an indigenous group was included, the most recent reference(s) is listed in this table. References for all studies are cited within the text of the results, and specific results from these studies and others can be referred to in the Appendix. References are included based on year of data, not year of publication.

[†]The data on TB burden among indigenous groups were abstracted from the most recent studies on indigenous groups, and correspond to the sources cited in this table. When more than one study was from the same period of time, the most recent TB data are presented as a range.

[‡]For most studies, uncertainty estimates were not available. When 95% confidence intervals were reported, they have been listed in the Appendix.

[§]Data for comparative TB burden were abstracted either from the same studies from which data were reported for the respective indigenous groups or from WHO reports. Additional information about these studies, including their source and scope, can be found in the Appendix.

[¶]Date study published; no date when data were collected was provided.

[#]The Pacific Islanders in the United States are recognized as an indigenous group, regardless of birthplace. Data segregated by birthplace were not available for this review. Data were not available that separately reported the TB burden in Native Hawaiians and Pacific Islanders in the United States.

^{***}Data reported for the Metis include only those self-registering as Metis. Anyone with at least some indigenous ancestry can register as Metis.

^{†††}As in the United States, the Pacific Islanders of New Zealand are considered an indigenous people, regardless of their birthplace, and as such, are incorporated into this review. TB = tuberculosis; WHO = World Health Organization; CDC = US Centers for Disease Control and Prevention.

The TB burden among indigenous peoples of the WHO African Region, as provided by all sources included in the systematic literature review ^{*}

Table A.1

Study, year	Country	Indigenous population	Type of study	Scope	Period	Age	TB burden			
							Cases (sample)	Prevalence/100000 [†]	Comparison prevalence/100000 Ratio [‡]	
Catra et al., 1997 ¹⁴	Mali	Peul and Dogon	Cross-sectional	Village/town/reserve	1997 [§]	Adults	4 (231)	1800	326 [¶]	5.5
Schelling et al., 2005 ¹⁵	Chad	Nomadic Fulani	Cross-sectional	Village/town/reserve	1999–2000	Adults	NA	4600 [¶]	429 [¶]	10.7

^{*}The TB burden reported is largely pulmonary. If a distinction was made in the studies between TB types, that has been noted in the table.

[†]TB cases not confirmed.

[‡]Unless otherwise noted, ratio is calculated as the listed prevalence or incidence of the indigenous group divided by the listed TB burden in the comparison group.

[§]Year of data collection not provided. Year of publication was used as substitute.

[¶]Nationwide statistic, retrieved from the WHO. Comparative statistics were for the same year as the data collection unless otherwise noted. NB: WHO prevalence measurements are estimates.

TB = tuberculosis; WHO = World Health Organization; NA = not available.

Table A.2

TB burden among indigenous peoples of the WHO Americas Region North America, as provided by all sources included in the systematic literature review *

Study, year	Country, province	Indigenous population	Type of study	Scope	Period	Age years	TB burden			Ratio [§]	
							Cases (sample)	Incidence/100000 [†]	Comparison incidence/100000 [‡]		
Blackwood et al., 2003 ⁴⁵	Canada, Manitoba	General	Cohort	Province	1992–1999	All	281 (–)	48.4	3.0 [¶]	16.1	
Bloss et al., 2011 ⁴⁴	USA, national	AI/AN	Cohort	National	2003–2008	All	914	5.9	1.1 [#]	5.4	
		NH/PI					362	14.7			13.4
Bracekner and Ellis, 1992 ⁴⁶	Canada, national	General	Cohort	National	1980–1990	All	NA	NA	NA	8 First Nations 9 Inuit	
Breault and Hoffman, 1997 ⁴⁷	USA, South Dakota	Sioux (Oglala)	Cohort	Village/town/reserve	1985–1994	All	77 (12 048)	63.9	2.8 [¶]	22.0	
CDC, 1991 ⁶⁴	USA, North and South Dakota	Sioux	Cross-sectional (outbreak investigation)	Village/town/reserve	1990 (January–May)	All	2 (7958)	25.1 ^{***}	10.3 ^{‡‡}	2.4	
CDC, 2010 ⁶	USA, national	AI/AN	Cohort	National	2008	All	137	5.9	1.1 [#]	5.4	
		NH/PI			2009		102	4.3	0.9 [#]	4.8	
				2008		69	15.9	1.1 [#]	14.5		
				2009		80	18.1	0.9 [#]	20.1		
CDC, 2011 ¹⁶	USA, national	AI/AN	Cohort	National	2010	All	146 (–)	6.1	0.9 [#]	6.8	
		NH/PI					94 (–)	20.6	0.9 [#]	22.9	
Clark et al., 2002 ⁴⁸	Canada, national	First Nations on-reserve	Cohort	National	1997–1999	All	NA	52.2	6.6 [#] All	7.9	
		British Columbia						32.3		4.9	
		Alberta						76.2		11.5	
		Saskatchewan						104.3		15.8	
		Manitoba						74.0		11.2	
		Ontario						15.9		2.4	
		New Brunswick and Nova Scotia						18.1		2.7	
		General	Cohort	Province	1990–2000			147 (–)	7.27 (95%CI 4.90–10.26)	0.18 [¶]	40.4
		Registered Indians	Cohort	Province	1986–2001	>20			93.8–95	4.4–5 [¶]	18.7–19
		With diabetes									
		Male						20 (1862)	NA		NA
		Female						50 (2802)	106.9		21.4
		Without diabetes									

Study, year	Country, province	Indigenous population	Type of study	Scope	Period	Age years	TB burden			Ratio [§]
							Cases (sample)	Incidence/ 100000 [†]	Comparison incidence/ 100000 [‡]	
Ellis et al., 2010 ¹⁷	Canada, national	Male	Cohort	National	2009	All	276 (17 603)	93.8	1.0 [#]	NA
		Female					245 (17 699)	NA		18.7
		Aboriginal (general)					342 (–)	27.8		27.8
		First Nations (North American Indian)					228 (3)	27.4		27.4
		Inuit					89 (–)	155.8		155.8
Enarson, 1998 ⁵¹	Canada, national	Metis	Cohort	National	1970–1995	All	NA	7.3	See graph, p S17	7.3
		General								>10
		AN		Province	1987–1994		NA	23.0		4.0
		General		Province	1994–1998		182 (750)	34.4	5.8 [¶]	19.6
		Inuit		Village/town/reserve	1990–2000	All	51 (–)	38–431	See figure, p 1354	310
Gessner, 1998 ⁵²	USA, Alaska	General indigenous	Cohort	Village/town/reserve	1986–2004	All	NA			29.2–331
		General	Cohort	National	1970–2001 (2001 rates listed)	<15	(–) 6683	15.3	1.3 [#]	25.2
		General	Cohort	Facility	1970–2001	All	9465 (82764 cases)	NA	0.6 [#]	1.43–RR CNS compared to PTB: aboriginal vs. non-aboriginal
		AI/AN	Cohort	National	1993–2002	All	(15 075 cases)	8.4	1.5 [#]	5.6
		All aboriginal	Cohort	National	2001	All	315 (587 cases nationwide)	NA	NA	18% of all cases were in the aboriginal population, versus 16% of all cases occurring in the non-aboriginal Canadian-born population [*]
Sheardown and Phippers, 2003 ⁶⁰	Canada, national	Status Indian	Case Control	City	1992–1996	All	200 (587)	112	1.5 [¶] 996 (See figure, p 153)	74.6
		Non-status Indian/Metis					63 (587)			11.3
		Inuit					52 (587)			47.3
		Aboriginals, on reserve					202 aboriginal cases			24
		Aboriginals, off reserve					45 (124 cases)			26.8
Wang et al., 2000 ⁶¹	Canada, British Columbia	Status Indians	Cohort	Province	1990–2004	<15	See map, p 92 and table, p 96	10.7	0.4 [¶]	4.0
		Alaska Natives Yukon								0.3
		Northwest Territories								0.5
		Nunavut								2.4
		Greenland								17.1
Yip et al., 2007 ⁶²	Canada, Alberta	Alaska Natives Yukon	Cohort	National	2004	All	See map, p 92 and table, p 96	25.8	6.5 [¶]	4.0
		Alaska Natives Yukon								0.3
		Northwest Territories								0.5
		Nunavut								2.4
		Greenland								17.1
Young, 2008 ⁶³	Far North (Canada, USA–Alaska)	Alaska Natives Yukon	Cohort	National	2004	All	See map, p 92 and table, p 96	12.9	44.5 [¶]	0.3
		Alaska Natives Yukon								0.5
		Northwest Territories								2.4
		Nunavut								17.1
		Greenland								17.1

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

* TB burden reported is largely pulmonary. If a distinction was made in the studies between TB types, that has been noted in the table.

[†] All incidence measurements are reported per year. If the study occurred over multiple years, the statistic reported is an average annual incidence, unless otherwise noted.

[‡] The comparison groups used in studies from Canada and the United States are the non-aboriginal native-born Canadians and White, non-Hispanic residents for the United States, unless otherwise noted.

[§] Unless otherwise noted, ratio is calculated as the listed incidence of the indigenous group divided by the listed TB burden in the comparison group.

[¶] State, district or provincial statistic, provided by author. Comparative statistics were for the same year as the data collection unless otherwise noted.

[#] Nationwide statistic, provided by author. Comparative statistics were for the same year as the data collection unless otherwise noted.

^{**} Authors of this systematic review calculated the incidence based on numbers provided in the designated article.

^{††} The comparison group is the TB incidence in the entire US population in 1990 as TB data was not disaggregated by race/ethnicity in the United States until 1993.

TB = tuberculosis; WHO = World Health Organization; CDC = US Centers for Disease Control and Prevention; NA = not available; CI = confidence interval; AI = American Indians; AN = Alaskan Natives; NH = Native Hawaiians; PI = Pacific Islanders; CNS = central nervous system manifestations of TB; PTB = pulmonary TB.

TB burden among indigenous peoples of the WHO Americas Region, Latin America and the Caribbean subdivision, as provided by all sources included in the systematic literature review*

Study, year	Country, province	Indigenous population	Type of study	Scope	Period	Age years	Cases (sample)	TB burden		
								Prevalence or incidence/100000 [†]	Comparison prevalence or incidence/100000 [‡]	Ratio [§]
Banzzi et al., 2001 ⁶⁶	Brazil, province not specified	Panara	Cross- sectional	Village/town/reserve	1995	All	10 (181)	Prevalence: 5525	Prevalence: 92 [¶]	60.1
Basta et al., 2004 ⁶⁷	Brazil, Rondoni	Surui	Cohort	City	2002	All	20 (864)	Incidence: 1388.9 (smear-positive) Incidence: 2314.8 (all forms)	Incidence: 43 [¶] /999 Incidence: 62 [¶] /999	32.3 37.3
Basta et al., 2006 ¹⁹	Brazil, Rondoni	Surui	Cross- sectional	City	2003	All	6 (920)	Prevalence: 815.2	Prevalence: 37.5 [#]	21.7
Basta et al., 2010 ²²	Brazil, Mato Grosso	Xavante	Cross- sectional	Village/town/reserve	2006	1	1 (476)	Prevalence: 210.8 ^{††}	Prevalence: 55 [¶]	3.8
			Cohort		1999–2004		37 (~560)	Incidence: 1289.6	Incidence: 48.5 [#]	26.6
Boia et al., 2009 ⁶⁸	Brazil-Amazonas	Arawak, Tukano, and Maku families (Arapaso, Baniwa, Desana, Kubeo, Hupda, Piratapuya, Tuyuca, Wanana, Tariana, and Tukano)	Cross- sectional	Village/town/reserve	2001	All	6 (333)	Prevalence: 1801.8 ^{††}	Prevalence: 75 [¶]	24.0
Buchillet and Gazin, 1998 ⁶⁹	Brazil, Amazonas	Arawak, Tukano, and Maku families (22 tribes)	Cohort	Facility	1990–1996 1977–1994	All	296 (23 000) 403 (2500)	Incidence: 200 Incidence: 920	Incidence: 50 [¶] Incidence: 73 [¶]	4.0 12.6
Culqui et al., 2010 ²¹	Peru, nationwide	Amazonic Ashaninka Shipibo Matsigenka Aguaruna Shawi Huambisa Yanesha Quichua-Amazonian Achuar Cashibo-Cacataybo Nomastshiguenga Quechua- Lamistas Anarakaire Harakmbut Shapra Aymara-Andes Quechua-Andes	Cohort, interviews	National	2008	All	68 38 23 20 13 11 7 6 4 6 2 2 1 1 1 84 417	Incidence: 89.0 Incidence: 230.0 Incidence: 226.0 Incidence: 42.3 Incidence: 92.0 Incidence: 106.0 Incidence: 153.1 Incidence: 197.3 Incidence: 121.4 Incidence: 400.9 Incidence: 53.9 Incidence: 64.0 Incidence: 133.0 Incidence: 360.5 Incidence: 45.0 Incidence: 175.1 Incidence: 103.7	Incidence: 106 ^{**}	0.8 2.2 2.1 0.4 0.8 1.0 1.4 1.9 1.2 3.8 0.5 0.6 1.3 3.4 0.4 1.7 1.0

Study, year	Country, province	Indigenous population	Type of study	Scope	Period	Age years	TB burden			Ratio [§]
							Cases (sample)	Prevalence or incidence/100000 [‡]	Comparison prevalence or incidence/100000 [‡]	
Escobar et al., 2001 ⁷⁰	Brazil, Rondoni	Indigenous, not specified	Cohort	Province	1992, 1994–1998	All	362 (NA)	Incidence: 1000	Incidence: 100 [#]	10.0
Fernandez de Larrea et al., 2002 ²⁶	Venezuela, Delta Amacuro	Warao People	Cross- sectional	Village/town/reserve	1999	<15	16 (502)	Prevalence: 320	Prevalence: 48 [#]	6.7
Garnelo et al., 2005 ^{71,‡‡}	Brazil, Special Indigenous Sanitary Districts	Indigenous, not specified, represented by district	Cohort, GIS	National	2000–2002	All	NA	Incidence: 302.1–1448.6	NA	>1000
Hurtado et al., 2003 ¹⁸	Paraguay	Ache natives	Cohort	Facility	1987–2002	All	80 (1000)	Incidence: 3700	Incidence: 49 [#]	75.5
Levino and de Oliveira, 2007 ⁷²	Brazil-Amazonas	Icama-Aiari	Cohort	District	2002	All	20 (32 180)	Incidence: 390	Incidence: 57 [#]	6.8
		Waupes-Papuri					16 (32 180)	Incidence: 200.4		3.5
		Tiquie					6 (32 180)	Incidence: 123.1		2.2
		Yanomami					32 (32 180)	Incidence: 2133.3		37.4
		Rio Negro					9 (32 180)	Incidence: 163.0		2.9
Machado Filho, 2008 ²⁰	Brazil-Amazonas	Indigenous, not specified	Cohort	Facility	1994–1998	All	322 (27 000)	Incidence: 239	Incidence: 64 [#]	3.7
					1999–2003		446 (31 407)	Incidence: 284	Incidence: 55 [#]	5.2
Marques et al., 2010 ²³	Brazil, Mato Grosso do Sul	Indigenous, not specified	Cohort	Province	2000	<15	224 (1096)	Incidence: 300	Incidence: 60 [#]	5.0
					2002			Incidence: 70	Incidence: 57 [#]	1.2
					2003			Incidence: 100	Incidence: 55 [#]	1.8
					2004			Incidence: 100	Incidence: 53 [#]	1.9
					2005			Incidence: 50	Incidence: 51 [#]	1.0
					2006			Incidence: 50	Incidence: 50 [#]	1.0
Romero-Amaro et al., 2008 ^{25,§§}	Venezuela, Zulia	Indigenous, including Anu, Bari, Wayuu, Yukpa	Cohort	Province	1996–2005	All	468 (889)	52% of TB cases occurred among the indigenous peoples	Incidence: 23.2 [#]	>1
Romero-Sandoval et al., 2007 ²⁴	Ecuador, Cotopaxi Province	Indigenous, not specified	Cross- sectional	Village/town/reserve	2001	All	44 (653)	Prevalence: 6700	Prevalence: 161 [#]	41.6
Sousa et al., 1997 ⁷³	Brazil-Amazonas	Yanomami	Cross- sectional	Village/town/reserve	1992	All	40 (625)	Prevalence: 6400 ^{‡‡} (all forms)	Prevalence: 68 [#]	94.1
van Crevel et al., 2004 ⁷⁴	Brazil-Amazonas, Surinam	Trio Indians	Cross- sectional; interview	Village/town/reserve	1995–2000	All	25 (–)	Prevalence: 420 (95%CI 270–610)	Prevalence: 60 [#]	7.0

^{*}The TB burden reported is largely pulmonary. If a distinction was made in the studies between TB types, that has been noted in the table.

[‡]All incidence measurements are reported per year. If the study occurred over multiple years, the statistic reported is an average annual incidence, unless otherwise noted.

^{‡‡}The comparison groups used in studies differed widely between studies. Numbers provided by authors for regional estimates were sometimes regional averages, while other times they were non-indigenous TB rates within those regions. Please refer to the specific studies for more details.

[§]Unless otherwise noted, ratio is calculated as the listed prevalence or incidence of the indigenous group divided by the listed TB burden in the comparison group.

^{##}Nationwide statistic, retrieved from the WHO. Comparative statistics were for the same year as the data collection unless otherwise noted. NB: WHO prevalence measurements are estimates.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

[#]State, district or provincial statistic, provided by author. Comparative statistics were for the same year as the data collection unless otherwise noted.

^{**}Nationwide statistic, provided by author. Comparative statistics were for the same year as the data collection unless otherwise noted.

^{††}Authors of this systematic review calculated the incidence or prevalence based on provided numbers in the designated article.

^{‡‡}This study presented data primarily in map format, revealing the large disparity in TB burden between regions with many indigenous and non-indigenous peoples. Specific incidence rates were only provided for the seven ‘Special Indigenous Sanitary Districts’ that had the highest incidence rates, all of which occurred within Legal Amazonia. Please refer to the study to obtain more specific data.

^{§§}This study did not provide a denominator for the indigenous population. However, 56% of TB cases (468/889) reported in Zulia, Venezuela, occurred among the indigenous population, with 89% of these cases occurring in the Wayuu peoples. The author concludes that indigenous are disproportionately affected by TB in comparison to the non-indigenous.

TB = tuberculosis; WHO = World Health Organization; GIS = Geographic Information System; NA = not available; CI = confidence interval.

TB burden among indigenous peoples of the WHO Eastern Mediterranean Region, as provided by all sources included in the systematic literature review^{*}

Table A.4

Study, year	Country, province	Indigenous population	Type of study	Scope	Period	Age years	Cases (sample)	TB burden		
								Incidence/100000 [†]	Comparison Incidence/100000	Ratio [‡]
Greene et al., 1992 ²⁷	Israel, The Negev Desert	Bedouins	Cohort	Facility	1987	All	11 (-)	18.1	8.7 [§] 1990	2.1

^{*}TB burden reported is pulmonary.

[†]All incidence measurements are reported per year.

[‡]Unless otherwise noted, ratio is calculated as the listed prevalence or incidence of the indigenous group divided by the listed TB burden in the comparison group.

[§]Nationwide statistic, retrieved from the WHO. NB: WHO prevalence measurements are estimates.

TB = tuberculosis; WHO = World Health Organization.

Table A.5

TB burden among indigenous peoples of the WHO European Region, as provided by all sources included in the systematic literature review ^{*}

Study, year	Country, province	Indigenous population	Type of study	Scope	Period	Age years	TB burden			Ratio [§]
							Cases (sample)	Prevalence or incidence/100000 [‡]	Comparison prevalence or incidence/100000 [‡]	
Casals et al., 2011 ³⁰	Spain	Roma	Cohort	Village/town/reserve	1985–2008	All	7 (380)	Incidence: 90.9 (95%CI 68.5–113.2)	Incidence: 17.0 [¶]	5.3
Chernukha et al., 2003 ⁷⁷	Russia, Far North	Indigenous, not specified	Cohort	Province	2003 [#]	All	NA	NA	NA	10.0 (Incidence)
Galkin and Vasilev, 1994 ⁷⁸	Russia, Far North	Indigenous	Cohort	Village/town/reserve	1988	All	NA	Incidence: 274.8	Incidence: 32.7 ^{**}	8.4
					1991		NA	Incidence: 176.1	Incidence: 25.6 ^{**}	6.9
					1992		NA	Incidence: 220.2	Incidence: 26.4 ^{**}	8.3
					1994	All	NA	Incidence: 303.7	Incidence: 68.4 ^{**}	4.4
Kucherov and Makarov, 2002 ²⁸	Russia, Chukotka Autonomous District	Chukotka Autonomous Indigenous	Cohort	District	1999		NA	Prevalence: 1159.5	Prevalence: 272.0 ^{**}	4.3
								Incidence: 85.2	Incidence: 37.6 ^{**}	2.3
								Prevalence: 608.0	Prevalence: 193.3 ^{**}	3.1
Ladefoged et al., 2011 ²⁹	Greenland	Inuit	Case control	National	2004–2006	All	146 (590 controls)	NA	Prevalence: 9.6 ^{¶¶}	15.3
Thomsen et al., 2004 ⁷⁶	Greenland	Inuit	Cohort	National	1990	All	NA	Prevalence: 85	Prevalence: 9.9 ^{¶¶}	8.6
					2001		NA	Prevalence: 185	Prevalence: 13.0 ^{¶¶}	14.2
Tyrylтин, 1990 ⁷⁹	Russia, Far North	Indigenous, not specified	Cohort	Province	1983–1987	All	NA	Incidence: 89.7	Prevalence: 41.9 ^{**}	2.1
Vasil'ev et al., 1992 ⁸⁰	Russia, Far North	Indigenous, not specified	Cohort	Province	1989	All	NA	Prevalence: 280	Prevalence: 220 [¶] /990	1.3
					1990		NA	Prevalence: 199.8		0.9

^{*} TB burden reported is largely pulmonary. If a distinction was made in the studies between TB types, that has been noted in the table.

[‡] All incidence measurements are reported per year. If the study occurred over multiple years, the statistic reported is an average annual incidence, unless otherwise noted.

[‡] The comparison groups used in studies differed widely between studies. Numbers provided by authors for regional estimates were sometimes regional averages, while other times they were non-indigenous TB rates within those regions. Please refer to the specific studies for more details.

[§] Unless otherwise noted, ratio is calculated as the listed prevalence or incidence of the indigenous group divided by the listed TB burden in the comparison group.

[¶] Nationwide statistic, retrieved from the WHO. Comparative statistics were for the same year as the data collection unless otherwise noted. NB: WHO prevalence measurements are estimates.

[#] Year of data collection not provided. Year of publication was used as substitute.

^{**} State, district, or provincial statistic, provided by the author. Comparative statistics were for the same year as the data collection unless otherwise noted.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

^{7/7} For Greenland, the comparative prevalence numbers are those provided by the WHO for Denmark.
TB = tuberculosis; WHO = World Health Organization; CI = confidence interval; NA = not available.

Table A.6

TB burden among indigenous peoples of the WHO South-East Asian Region, as provided by all sources included in the systematic literature review*

Author, year	Country, province	Indigenous population	Type of study	Scope	Period	Age years	TB burden			
							Cases (sample)	Prevalence or incidence/100000 [†]	Comparison prevalence or incidence/100000 [‡]	Ratio [§]
Balasubramanian et al., 1995 ⁸⁴	India, Tamil Nadu	Jawadhu-Hills tribes	Cross- sectional	Village/town/reserve	1992–1993	15	12 (5755)	Prevalence: 209 [¶]	Prevalence: 461 [#]	0.5
Bhat et al., 2009 ⁸¹	India, Madhya Pradesh	Tribals, not specified (refugees)	Cross- sectional	District	2007–2008	15	– (22 270)	Prevalence: 387 (95% CI 272–502)	Prevalence: 299 ^{**} 2006	1.3
Bhatia et al., 2002 ⁴²	India, Himachal Pradesh	Tibetans (refugees)	Cross- sectional	Village/town/reserve	1996	All	1575 (53 959)	Incidence: 770	Incidence: 216 [#]	3.6
CDC, 2005 ¹⁰⁴	Thailand, refugee camp	Hmong (refugees)		Village/town/reserve	2004–2006	All	272 (15 455)	Prevalence: 1760 ^{¶¶}	Prevalence: 193 ^{**}	9.1
Chakma et al., 1996 ⁸²	India, Madhya Pradesh	Saharia	Cross- sectional	Village/town/reserve	1991–1992	15	96 (22 250)	Prevalence: 1500	Prevalence: 970 ^{**}	1.6
Datta et al., 2001 ³¹	India, Tamil Nadu	Jawadhu-Hills tribes	Cross- sectional	Village/town/reserve	1989	All	– (16 017)	Prevalence: 840	Prevalence: 459 [#]	1.8
Kaulagekar and Radkar, 2007 ⁸⁵	India, national	Scheduled tribes	Cohort	National	1998–1999	All	NA	Prevalence: Male 107	Prevalence: Male 74 ^{††}	1.5
								Prevalence: Female 263	Prevalence: Female 54 ^{††}	4.9
Kerketta et al., 2009 ⁴⁰	India, Orissa	Forest peoples, general	Cross- sectional	Village/town/reserve	2009 ^{†††}	Elderly (μ = 68)	15 (312)	Prevalence: 4808 ^{†††}	Prevalence: 275 [#]	17.5
		Dongria Kondh					9 (50)	Prevalence: 18 000 ^{¶¶}		65.5
		Kutia Kondh					3 (112)	Prevalence: 2679 ^{¶¶}		9.7
		Langia Saora					0 (50)	Prevalence: 0 ^{¶¶}		0.0
		Pandi Bhuiyan					3 (100)	Prevalence: 3000 ^{¶¶}		10.9
Marras et al., 2003 ^{103,88,89,¶¶}	Canada, Toronto	Tibetans (refugees)	Cohort	Facility	1998–2000	Adults	24 (525)	Incidence: 4571	Incidence: Canada 6.4 [#]	714.2 (Canada)
									Incidence: India 216 [#]	21.2 (India)
Murhekar et al., 2004 ³⁹	India, Car Nicobar	Nicobarese	Cross- sectional	Province	2001–2002	15	– (10 570)	Prevalence: 735.3	Prevalence: 436 [#]	1.7
Narang et al., 1999 ³⁸	India, Maharashtra	Maharashtra, indigenous tribes (<i>n</i> = 46)	Cross- sectional	District	1989–1990	All	– (20 596)	Prevalence: 133	Prevalence: 144 ^{**}	0.9
		Gond					– (5574)	Prevalence: 136		0.9
		Gond gawari					– (5574)	Prevalence: 100		0.7
		Mana					– (150)	Prevalence: 730		5.1
		Pawara					– (170)	Prevalence: 612		4.3
		Raj Gond					– (5574)	Prevalence: 196		1.4
Nelson et al., 2005 ¹⁰¹	India, national	Tibetans (refugees)	Cross- sectional	National	1994–1996	All	1197 (47 491)	Incidence: 835	Incidence: 216 [#]	3.9
Oeltmann et al., 2008 ⁴¹	Thailand, refugee Camp	Hmong (refugees)	Cross- sectional	Village/town/reserve	2004–2005	All	272 (15 455)	Prevalence: 1760 ^{¶¶}	Prevalence: 193 [#]	9.1
Rao et al., 2010 ³⁶	India, Madhya Pradesh	Bharia	Cross- sectional	District	2008	15	6 (1390)	Prevalence: 432 (95% CI 121–742)	Prevalence: 294 [#]	1.5

Author, year	Country, province	Indigenous population	Type of study	Scope	Period	Age years	TB burden		
							Cases (sample)	Prevalence or incidence/100000 [†]	Comparison prevalence or incidence/100000 [‡] Ratio [§]
Rao et al., 2010 ³⁴	India, Madhya Pradesh	Tribals, not specified	Cross- sectional	District	2007–2008	15	83 (20 000)	Prevalence: 415 [¶]	Prevalence: 294 [#] 1.4
Rao et al., 2010 ³²	India, Madhya Pradesh	Saharia	Cross- sectional	District	2007–2008	All	166 (11 116)	Prevalence: 1518	Prevalence: 294 [#] 5.2
Rao et al., 2012 ³⁷	India, Madhya Pradesh	Saharia	Cross- sectional	District	2007–2008	15	113 (9538)	Prevalence: 1394 [¶]	Prevalence: 294 [#] 4.7
Sharma and Tiwari, 2007 ⁸³	India, Madhya Pradesh	Saharia	Cross- sectional	Village/town/reserve	2005–2006	All	164 (355)	Prevalence: 46 197 [¶]	Prevalence: 335 [#] 137.9
Sharma et al., 2010 ³⁵	India, Madhya Pradesh	Bhil	Cross- sectional	Village/town/reserve	2006–2007	All	4 (169)	Prevalence: 2366 [¶]	Prevalence: 314 [#] 7.5
Truong et al., 1997 ^{102,§§,¶¶}	United States, Minnesota	Tibetans (refugees)	Cross- sectional	Facility	1992–1994	All	16 (191)	Incidence: 8377 (2–3 years)	Incidence: USA 11 [#] 761.5 (US) 38.8 (India)
		Hmong (refugees)					39 (2456)	Incidence: 1588 (2–3 years)	Incidence: India 216 [#] 144.4 (US)
Yadav et al., 2010 ³³	India, Madhya Pradesh	Baiga	Cross- sectional	Village/town/reserve	2008	All	2 (2359)	Prevalence: 146 (95%CI 0–318)	Incidence: Thailand 137 [#] 11.6 (Thai) 0.5

^{*}The TB burden reported is for pulmonary TB.

[†]All incidence measurements are reported per year. If the study occurred over multiple years, the statistic reported is an average annual incidence, unless otherwise noted.

[‡]The comparison groups used in studies differed widely between studies. Numbers provided by authors for regional estimates were sometimes regional averages, while at other times they were non-indigenous TB rates within those regions. Please refer to the specific studies for more details.

[§]Unless otherwise noted, ratio is calculated as the listed prevalence or incidence of the indigenous group divided by the listed TB burden in the comparison group.

[¶]Authors of this systematic review calculated the incidence or prevalence based on provided numbers in the designated article.

[#]Nationwide statistic, retrieved from the WHO. Comparative statistics were for the same year as the data collection unless otherwise noted. NB: WHO prevalence measurements are estimates.

^{**}State, district, or provincial statistic, provided by author. Comparative statistics were for the same year as the data collection unless otherwise noted.

^{††}Nationwide statistic, provided by the author. Comparative statistics were for the same year as the data collection unless otherwise noted.

^{†††}Year of data collection not provided. Year of publication was used as substitute.

^{§§}Authors of this study referred to their measurements as prevalence. However, our understanding of their study has led us to believe they are actually incidence measurements, and as such, we have reported them as incidence.

^{¶¶}The incidence measurement for this study is not the average annual incidence. It is for the entire study period (1992–1993), or 2–3 years.

TB = tuberculosis; WHO = World Health Organization; NA = not available; CI = confidence interval; CDC = Centers for Disease Control and Prevention.

Table A.7

TB burden among indigenous peoples of the WHO Western Pacific Region, as provided by all sources included in the systematic literature review ^{*}

Study, year	Country, province	Indigenous population	Type of study	Scope	Period	Age years	TB burden			
							Cases (sample)	Prevalence or incidence/100000 [†]	Comparison prevalence or incidence/100000 [‡]	Ratio [§]
Barry and Konstantinos, 2009 ⁴	Australia, national	Aborigines/Torres Strait Islanders	Cohort	National	2007	All	35 (–)	Incidence: 6.6	Incidence: 0.9 [¶]	7.3
	Capital Territory						0 (–)	Incidence: 0	Incidence: 0.9 [#]	0.0
	New South Wales			—			3 (–)	Incidence: 1.9	Incidence: 1.2 [#]	1.6
	Northern Territory						21 (–)	Incidence: 32.2	Incidence: 2.4 [#]	13.4
	Queensland						7 (–)	Incidence: 4.7	Incidence: 0.4 [#]	11.8
	Southern Australia						3 (–)	Incidence: 10.5	Incidence: 0.8 [#]	13.1
	Tasmania						0 (–)	Incidence: 0	Incidence: 0.5 [#]	0.0
	Victoria						1 (–)	Incidence: 2.9	Incidence: 0.9 [#]	3.2
	Western Australia						0 (–)	Incidence: 0	Incidence: 0.8 [#]	0.0
	New Zealand, Auckland	Maori	Cohort	City	1992–1993	All	24 (–)	Incidence: 37.8	Incidence: 2.7 [#]	14.0
Calder and Priest, 2006 ⁹¹		Pacific Islanders					53 (–)	Incidence: 70.9	Incidence: 2.7 [#]	26.3
	Taiwan, national	Aborigines	Cohort	National	1996–2003	<20	NA	Incidence: 81.5	Incidence: 9.6 [¶]	8.5
	Taiwan, eastern	Aborigines	Case control	Village/town/reserve	2000–2006	All	60 (–)	Incidence: 321.1 (all) 152.1 (smear +)	Incidence: 66.7 [¶]	2.3–4.8
	Taiwan, national	Aborigines	Cohort	National	1996–2006	All	NA	Incidence: 176	Incidence: 65 [¶]	2.7
	New Zealand, national	Maori	Cohort	National	2000–2004	All	332 (526 281)	Incidence: 20.46	Incidence: 1.43 [¶]	14.3
					1995–1999		352 (523 371)	Incidence: 23.55	Incidence: 2.56 [¶]	9.2
					2000–2004		359 (200 253)	Incidence: 48.79	Incidence: 1.43 [¶]	34.1
		Pacific Islanders			1995–1999		303 (173 181)	Incidence: 51.77	Incidence: 2.56 [¶]	20.2
	New Zealand, national	Maori	Cohort	National	1992–2001	<16	NA	Incidence: 6.4	Incidence: 0.6 [¶]	10.7
		Pacific Islanders					NA	Incidence: 15.2		25.3
Hsu et al., 1990 ⁹⁸	China, Northeast	E-Lun Chun (Ortochen)	Cohort	Province	1985	All	NA	Incidence: 6500	Incidence: 153 ^{**¶} 990	42.5
	Taiwan	Aborigines	Cohort	National	2002	All	NA	Incidence: 289.8	Incidence: 74.6 [¶]	3.9
	Northern Territory	Aborigines/Torres Strait Islanders descendants	Cohort	National	2003	All	38 (–)	Incidence: 8.7	Incidence: 0.9 [¶]	9.7
	Capital Territory						0 (–)	Incidence: 0	Incidence: 2.0 [#]	0.0
Howie et al., 2005 ⁹²										
Hsu et al., 1990 ⁹⁸										
Hsueh et al., 2006 ⁹⁶										
Li et al., 2004 ⁸⁶										

Study, year	Country, province	Indigenous population	Type of study	Scope	Period	Age years	TB burden		
							Cases (sample)	Prevalence or incidence/100000 [†]	Comparison or prevalence or incidence/100000 [‡] Ratio [§]
Miller et al., 2002 ⁸⁷	Australia, national						5 (–)	Incidence: 4.1	Incidence: 0.8 [#] 5.1
	New South Wales						20 (–)	Incidence: 34.9	Incidence: 1.8 [#] 19.4
	Queensland						6 (–)	Incidence: 4.9	Incidence: 0.8 [#] 6.1
	Southern Australia						2 (–)	Incidence: 8.1	Incidence: 0.8 [#] 10.1
	Tasmania						0 (–)	Incidence: 0	Incidence: 0.5 [#] 0.0
	Victoria						0 (–)	Incidence: 0	Incidence: 1.0 [#] 0.0
	Western Australia						5 (–)	Incidence: 8	Incidence: 0.3 [#] 26.7
	Australia, national	Aborigines/Torres Strait Islanders descendants	Cohort	National	2001	All	42 (–)	Incidence: 9.8	Incidence: 1 [¶] 9.8
	Capital Territory						0 (–)	Incidence: 0	Incidence: 0.4 [#] 0.0
	New South Wales						1 (–)	Incidence: 0.8	Incidence: 1.4 [#] 0.6
	Northern Territory						25 (–)	Incidence: 44.4	Incidence: 2.7 [#] 16.4
	Queensland						9 (–)	Incidence: 7.6	Incidence: 0.9 [#] 8.4
	Southern Australia						1 (–)	Incidence: 4.1	Incidence: 0.6 [#] 6.8
	Tasmania						0 (–)	Incidence: 0	Incidence: 1.7 [#] 0.0
	Victoria						3 (–)	Incidence: 12.2	Incidence: 1 [#] 12.2
	Western Australia						3 (–)	Incidence: 4.9	Incidence: 0.7 [#] 7.0
	Australia, national	Aborigines/Torres Strait Islanders descendants	Cohort	National	1994	All	199 (–)	Incidence: 10.6	Incidence: 6.7 ^{**} 1.6
	Australia, Far North Queensland	Aborigines, Papua New Guineans	Cohort	Province	1998–2002	All	6692 cases	NA	NA
	Australia, Far North Queensland	Aborigines	Cohort	Province	1993–1997	All	5787 cases	Incidence: 35.9	Incidence: 2.3 [#] 15.6
	New Zealand	Maori	Cohort	District	1993–1997	All	50 (27 818)	Incidence: 12.5	Incidence: 11 ^{**} 1.1
	China, national	Bai	Cohort	National	1992–2001	All	244 (317 751)	Prevalence: 47	Prevalence: 134 [¶] 0.4
Wang and You, 1994 ⁹⁹	Chaoxian (Chinese Korean)				1990	All	NA	Prevalence: 61	[50 (urban) 140 (rural)] 0.5
	Hui							Prevalence: 181	1.4
	Kazakh							Prevalence: 180	1.3
	Li							Prevalence: 262	2.0
	Manchu							Prevalence: 90	0.7
Miao (Hmong)								Prevalence: 163	1.2

Study, year	Country, province	Indigenous population	Type of study	Scope	Period	Age years	TB burden		
							Cases (sample)	Prevalence or incidence/100000 [†]	Comparison prevalence or incidence/100000 [‡] Ratio [§]
Wu et al., 2002 ¹⁰⁰	China, Sichuan	Mongols (Menggu)	Cohort	Province	2000	NA	NA	Prevalence: 170	1.3
		Tibetans						Prevalence: 106	0.8
		Tujia						Prevalence: 247	1.8
		Uyghur						Prevalence: 274	2.0
		Zhuang						Prevalence: 103	0.8
Yu et al., 2004 ⁹⁵	Taiwan	Ethnic minorities, not specified	Cohort	National	2001	All	520 (–)	Prevalence: 395	Prevalence: urban 172 [#] 2.3
		Aborigines							Prevalence: rural 126 [#] 3.1
								Incidence: 233.5	Incidence: 56.5 [¶] 4.1
								Incidence: 278.4	Incidence: 53.5 [¶] 7.1

^{*}The TB burden reported is largely pulmonary. If a distinction was made in the studies between TB types, that has been noted in the table.

[†]All incidence measurements are reported per year. If the study occurred over multiple years, the statistic reported is an average annual incidence, unless otherwise noted.

[‡]For Australia, New Zealand, and Taiwan, comparison rates provided by authors were for non-aboriginal citizens in those countries.

[§]Unless otherwise noted, ratio is calculated as the listed prevalence or incidence of the indigenous group divided by the listed TB burden in the comparison group.

[¶]Nationwide statistic, provided by the author. Comparative statistics were for the same year as the data collection unless otherwise noted.

[#]State, district or provincial statistic, provided by the author. Comparative statistics were for the same year as the data collection unless otherwise noted.

^{***}Nationwide statistic, retrieved from the WHO. Comparative statistics were for the same year as the data collection unless otherwise noted. NB: WHO prevalence measurements are estimates.

TB = tuberculosis; WHO = World Health Organization; NA = not available.